

New record of *Furipterus horrens* (Cuvier, 1828) (Mammalia, Chiroptera) from the Cerrado of Tocantins state with a compilation of the known distribution within Brazil

Roberto Leonan Morim Novaes*, Renan de França Souza, Saulo Felix, Cristal Sauwen, Gabriella Jacob and Leonardo dos Santos Avilla

Universidade Federal do Estado do Rio de Janeiro, Departamento de Zoologia, Laboratório de Mastozoologia. Av. Pasteur, 459, sala 501, IBIO. CEP 22290-240. Rio de Janeiro, RJ, Brazil.

* Corresponding author. E-mail: robertoleonan@gmail.com

ABSTRACT: *Furipterus horrens* is an insectivorous bat that occurs from Costa Rica to southeastern Bolivia and southern Brazil, with records in the Amazon, Caatinga, Cerrado and Atlantic Forest biomes. Despite this broad distributional range across South America, the species is currently known from few localities, and its habitat preferences are poorly known. We report the first record of *Furipterus horrens* for the Tocantins state, northern Brazil, based on four individuals collected in two caves surrounded by Neotropical savanna (Cerrado) in the Aurora do Tocantins municipality.

The family Furipteridae (Mammalia: Chiroptera) is currently composed by only two genera, both monotypic: *Amorphochilus schnablii* Peters, 1877 and *Furipterus horrens* (Cuvier, 1828) (Simmons 2005). *Furipterus horrens* occurs in Costa Rica, Panama and Trinidad southward to southeastern Bolivia and southern Brazil (Emmons 1990, Gardner 2008, Peracchi *et al.* 2011). Within Brazil this species has been recorded for 12 states in all regions, covering the Amazon, Caatinga, Cerrado and Atlantic Forest biomes (Reis and Gazarini 2007, Tavares *et al.* 2008, Peracchi *et al.* 2011). We report here the first record of *F. horrens* for the state of Tocantins, northern Brazil. This is the northernmost record within Brazilian Cerrado, being about 300 km away from the last record to the Cerrado.

The record of this species was made during the survey activities of the bat fauna in a karstic area of the Aurora do Tocantins municipality, Southeast Tocantins state, northern Brazil (Figure 1). The site has several massive limestones with dozens of caves (Figure 2). The massifs are inserted into a mosaic landscape composed of large remnants of the Neotropical savanna (Cerrado) and agricultural properties with cattle ranching and maize, cassava, rice and soybeans plantations. We performed eight nights of bats sampling, being five nights of 12 h (18:00 - 06:00 h) and three nights of 6 h (18:00 - 00:00h), using 10 mist nets (Zootech® 9 x 3 m, 20 mm), totalized 21,260 m².h of sampling effort (Straube and Bianconi 2002). The mist-nets were placed (1) at the caves entrance (6,390m².h), (2) to a distance of 15m from the caves entrance (6,380m².h), (3) in edges, trails and glades in the fragments of Cerrado (6,380m².h) and (4) around artificial ponds (2,110m².h). We also performed diurnal active searches and manual catches using hand-nets. Individuals collected under permits SISBIO/IBAMA (4028-1 / 28717) (MN78114, MN78116, MN78115, MN78117) were deposited in the mammal collection of the Museu Nacional do Rio de Janeiro, Rio de Janeiro, Brazil. We followed the guidelines of Sikes *et al.*

(2011) use of animals for scientific purposes.

In January 14, 2012 four adult individuals of *F. horrens* were collected at the Gruta dos Moura (12°34'53.4" S/ 46°30'59.0" W, altitude ca. 400 m), Aurora do Tocantins municipality, Tocantins state, northeastern Brazil. The specimens collected during the present study were part of a colony with more than 100 individuals sheltering in the ceiling at the cave entrance at about 1.50 m above the ground. Two adult individuals were males with abdominal testes (MN78114: forearm 34.1mm, weight 4.0g and MN78116: forearm 34.0, weight 3.0g) and two were inactive females (MN78115: forearm 34.5, weight 4.0g and MN78117: forearm 35.6, weight 3.5g) (Figure




FIGURE 1. Capture localities of *Furipterus horrens* within Brazil (blue circles), including the new record from Aurora do Tocantins (red circle) on Brazilian biomes.

3). Measurements and morphological characters fall within the expected variation for the species (e.g. Husson 1978, Brosset and Charles-Dominique 1990, Simmons and Voss 1998, Reis and Gazarini 2007, Peracchi *et al.* 2010). *F. horrens* can be easily identified due to its peculiar morphology, including a reduced thumb included in the wing membrane with a rudimentary nail, and a truncated snout with a disk-shaped extremity, the large and round-shaped ears funnel-shaped, with a short and triangular tragus (Vieira 1942, Cabrera and Yapes 1960).

Mist nets were not effective to capture this species, since even the grids placed in front of two previously identified shelters were not able to capture any individual. *F. horrens* has a diet composed exclusively by insects (Wilson 1973,



FIGURE 2. Massive limestones includes the Gruta dos Moura, in agricultural landscape composed of Cerrado fragments, cornfield and pasture.



FIGURE 3. *Furipterus horrens* female captured in a cave in the city of Aurora do Tocantins, southeastern of Tocantins state, Brazil.

Nowak 1994), located with the help of echolocation, which may facilitate the mist nets detection (Arita and Fenton 1997; Schnitzler and Kalko 2001), and appears to have high flight maneuverability. Accordingly, we observed several individuals quickly deviating from the mist-nets. Gregorin *et al.* (2008) during a survey of bats in the Piauí State, captured this species only manually in roosts. Within Brazil, *F. horrens* is now documented in 26 locations (Table 1, Figure 1) including the present record from Tocantins. The knowledge about the distribution of this species within Brazil is clearly incomplete, and probably biased to sampling methods that have been most commonly used in surveys.

TABLE 1. Occurrence locations of *Furipterus horrens* in Brazil.

LOCALITY	STATE	BIOME	COORDINATES	AUTHOR
80 km north of Manaus	AM	Amazon	02°30' S, 59°43' W	Voss and Emmons (1996)
Taperinha, Santarém	PA	Amazon	02°32' S, 54°17' W	Piccinini (1974)
Manaus	AM	Amazon	03°06' S, 60°00' W	Piccinini (1974)
Itapipoca	CE	Atlantic Forest	03°29' S, 39°34' W	Piccinini (1974)
Irauçuba	CE	Caatinga	03°44' S, 39°47' W	Fabián (2008)
PN Ubajara	CE	Caatinga	03°48' S, 40°42' W	Uieda <i>et al.</i> (1980)
Quixadá	CE	Caatinga	04°58' S, 39°01' W	Ástua and Guerra (2008)
Serra das Almas, Crateús	CE	Caatinga	05°10' S, 40°40' W	Silva <i>et al.</i> (2004)
FLONA Carajás	PA	Amazon	06°04' S, 50°13' W	Tavares <i>et al.</i> (2012)
Exu	PE	Caatinga	07°30' S, 39°42' W	Mares <i>et al.</i> (1981)
Brejo da Madre de Deus	PE	Caatinga	08°08' S, 36°22' W	Souza <i>et al.</i> 2004
Rio Formoso	PE	Atlantic Forest	08°39' S, 35°09' W	Bonato and Facure (2000)
PN Serra das Confusões	PI	Caatinga	09°13' S, 43°29' W	Gregorin <i>et al.</i> (2008)
Canindé de São Francisco	SE	Caatinga	09°38' S, 37°47' W	Ástua and Guerra (2008)
Aurora do Tocantins	TO	Cerrado	12°34' S, 46°30' W	Present study
Southern of Bahia	BA	Atlantic Forest	?	Faria <i>et al.</i> (2006)
Mambáí	GO	Cerrado	14°29' S, 46°06' W	Coimbra <i>et al.</i> (1982)
Nascentes Rio Vermelho	GO	Cerrado	14°24' S, 46°11' W	Esbérard <i>et al.</i> (2005)
PNM Cavernas do Peruaçu	MG	Cerrado	15°05' S, 44°15' W	Tavares <i>et al.</i> (2010)
Rural area	DF	Cerrado	15°44' S, 47°57' W	Bredt and Uieda (1996)
FLONA Rio Preto	ES	Atlantic Forest	18°21' S, 39°50' W	Duda <i>et al.</i> (2012)
APA Cairuçu, Parati	RJ	Atlantic Forest	23°17' S, 44°31' W	Pol <i>et al.</i> (2003)
PE Intervalles	SP	Atlantic Forest	24°16' S, 48°24' W	Portfors <i>et al.</i> (2000)
PETAR, Iporanga	SP	Atlantic Forest	24°35' S, 48°37' W	Trajano (1985)
ESEC Juréia-Itatins	SP	Atlantic Forest	24°40' S, 47°21' W	Gimenez and Ferrarezi (2004)
Colônia Hansa, Corupá	SC	Atlantic Forest	26°21' S, 48°29' W	Cherem <i>et al.</i> (2004)

ACKNOWLEDGMENTS: We are grateful to the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for supporting the project; Anselmo Rodrigues and Wagner José Moura for structural and logistical support on the performed activities; Municipal district of Aurora do Tocantins by the accommodation granting; and Valéria da Cunha Tavares for valuable suggestions and corrections to the original manuscript.

LITERATURE CITED

- Arita, H.T. and M.B. Fenton. 1997. Flight and echolocation in the ecology and evolution of bats. *Tree* 12(2): 53-58.
- Ástua, D. and D.Q. Guerra. 2008. Caatinga bats in the Mammal Collection of the Universidade Federal de Pernambuco. *Chiroptera Neotropical* 14(1): 326-338.
- Bernard, E., V.C. Tavares and E. Sampaio. 2011. Compilação atualizada das espécies de morcegos (Chiroptera) para a Amazônia Brasileira. *Biota Neotropica* 11(1): 1-13.
- Bonato, V. and K.G. Facure. 2000. Bat predation by the fringelipped bat *Trachops cirrhosus* (Chiroptera: Phyllostomidae). *Mammalia* 64: 241-243.
- Bredt, A. and W. Uieda. 1996. Bats from urban and rural environments of the Distrito Federal, mid-western Brazil. *Chiroptera Neotropical* 2(2): 54-57.
- Brosset, A. and P. Charles-Dominique. 1990. The bats from French Guiana: a taxonomic, faunistic and ecological approach. *Mammalia* 54(4): 509-560.
- Cabrera, A. and Yepes, J. 1960. *Mamíferos Sudamericanos*. Buenos Aires: Ediar. 238 p.
- Cherem, J.J., P.C. Simões-Lopes, S. Althoff and M.E. Graipel. 2004. Lista dos mamíferos do estado de Santa Catarina, sul do Brasil. *Mastozoología Neotropical* 11(2): 151-184.
- Coimbra, C.E.A., M.M. Borges, D.Q. Guerra and D.A. Mello. 1982. Contribuição à zoogeografia e ecologia dos morcegos em regiões de Cerrado do Brasil central. *Boletim Técnico do IBDF* 7: 33-38.
- Duda, R., J. Dalapiccola and L.P. Costa. 2012. First record of the smoky bat *Furipterus horrens* (F. Cuvier, 1828) (Mammalia: Chiroptera) in the state of Espírito Santo, southeastern Brazil. *Check List* 8(6): 1362-1364.
- Emmons, L.H. 1990. *Neotropical rainforest mammals: a field guide*. Chicago: University of Chicago Press. 281 p.
- Esbérard, C.E.L., J.A. Motta and C. Perigo. 2005. Morcegos cavernícolas da Área de Proteção Ambiental (APA) Nascentes do Rio Vermelho, Goiás. *Revista Brasileira de Zootecias* 7(2): 311-325.
- Fabián, M.E. 2008. Quirópteros do bioma Caatinga, no Ceará, Brasil, depositados no Museu de Ciências da Fundação Zoobotânica do Rio Grande do Sul. *Chiroptera Neotropical* 14(1): 354-359.
- Faria, D., B. Soares-Santos and E. Sampaio. 2006. Bats from the Atlantic rainforest of southern Bahia, Brazil. *Biota Neotropica* 6(2): 1-13.
- Gardner, A.L. 2008. Family Furipteridae; p. 389-392. In A.L. Gardner (ed.). *Mammals of South America*. Chicago: The University of Chicago Press. 669 p.
- Gimenez, E.A. and H. Ferrarezzi. 2004. Diversidade de morcegos no sudesta da Mata Atlântica; p. 214-330 In O.A.V. Marques and W. Duleba (ed.). *Estação Ecológica Juréia-Itatins: ambiente físico, flora e fauna*. Ribeirão Preto: Holos.
- Gregorin, R., A.P. Camignotto and A.R. Percequillo. 2008. Quirópteros do Parque Nacional da Serra das Confusões, Piauí, nordeste do Brasil. *Chiroptera Neotropical* 14: 366-383.
- Husson, A.M. 1978. The mammals of Surinam. *Zoölogische Monographien van het Rijksmuseum van Natuurlijke Historie* 2: 569 p.
- Mares, M.A., M.R. Willig, K.E. Streilein and T.E. Lacher. 1981. The mammals of northeastern Brazil: a preliminary assessment. *Annals of Carnegie Museum* 50: 81-137.
- Nowak, R.M. 1994. *Walker's bats of the world*. Baltimore: John Hopkins University Press. 287p.
- Peracchi, A.L., P.H. Gallo, D. Dias, I.P. Lima and N.R. Reis. 2010. Ordem Chiroptera; p. 293-461. In N.R. Reis, A.L. Peracchi, M.N. Fregonezi and B.K. Rossaneis (org.). *Mamíferos do Brasil: guia de identificação*. Rio de Janeiro: Technical Books Editora. 560 p.
- Peracchi, A.L., I.P. Lima, N.R. Reis, M.R. Nogueira and H. Ortêncio-Filho. 2011. Ordem Chiroptera; p. 155-234 In N.R. Reis, A.L. Peracchi, W.A. Pedro and I.P. Lima (ed.). *Mamíferos do Brasil*, 2ª Edição. Londrina: Editora da Universidade Estadual de Londrina.
- Piccinini, R.S. 1974. Lista provisória dos quirópteros da coleção do Museu Paraense Emílio Goeldi (Chiroptera). *Boletim do Museu Paraense Emílio Goeldi, Nova Série Zoologia* 77: 1-32.
- Pol, A., M.R. Nogueira and A.L. Peracchi. 2003. Primeiro registro da família Furipteridae (Mammalia, Chiroptera) para o Estado do Rio de Janeiro, Brasil. *Revista Brasileira de Zoologia* 20(3): 561-563.
- Portfors, C.V., M.B. Fenton, L.M.S. Aguiar, J.E. Baumgarten, M.J. Vonhof, S. Bouchard, D.M. Faria, W.A. Pedro, N.I.L. Rauntenbach and M. Zortéa. 2000. Bats from Fazenda Intervales, Southeastern Brazil: species account and comparison between different sampling methods. *Revista Brasileira de Zoologia* 17(2): 533-538.
- Reis, N.R. and J. Gazarini. 2007. Família Furipteridae, p. 137-138. In N.R. Reis, A.L. Peracchi, W.A. Pedro and I.P. Lima (ed.). *Morcegos do Brasil*. Londrina: Editora da Universidade Estadual de Londrina. 253 p.
- Schnitzler, H.U. and E.K.V. Kalko. 2001. Echolocation by insect-eating bats. *BioScience* 51(7): 557-569.
- Sikes, R.S., W.L. Gannon and Animal Care and Use Committee of the American Society of Mammalogists. 2011. Guidelines of the American Society of Mammalogists for the use of wild mammals in research. *Journal of Mammalogy* 92(1): 235-253.
- Silva, S.S.P., P.G. Guedes, A.R. Camardella and A.L. Peracchi. 2004. Survey of bats (Mammalia, Chiroptera), with comments on reproduction status, in Serra das Almas Private Heritage Reserve, in the state of Ceará, northeastern of Brazil. *Chiroptera Neotropical* 10: 191-195.
- Simmons, N.B. 2005. Order Chiroptera; p. 312-529 In D.E. Wilson and D.M. Reeder (ed.). *Mammals Species of the World: a taxonomic and geographic reference*. Chicago: Johns Hopkins University Press.
- Simmons, N.B. and R.S. Voss. 1998. The mammals of Paracou, French Guiana: a neotropical lowland rainforest fauna. Part 1. Bats. *Bulletin of the American Museum of Natural History* 237: 1-219.
- Souza M.A.N., A. Langguth and E.A. Gimenez. 2004. Mamíferos dos brejos de altitude da Paraíba e Pernambuco; p. 229-254. In K.C. Porto, J.J.P. Cabral and M. Tabarelli (ed.). *Brejos de altitude em Pernambuco e Paraíba: história natural, ecologia e conservação*. Brasília: Ministério do Meio Ambiente/PROBIO/CEPAN.
- Straube, F.C. and G.V. Bianconi. 2002. Sobre a grandeza e a unidade utilizada para estimar esforço de captura com utilização de redes-de-neblina. *Chiroptera Neotropical* 8: 150-152.
- Tavares, V.C., C.F.S. Palmuti, R. Gregorin and T.T. Dornas. 2012. Morcegos; p. 162-179. In F.D. Martins, A.F. Castilho, J. Campos, F.M. Hatano and S.G. Rolim. (org.). *Fauna da Floresta Nacional de Carajás: estudos sobre vertebrados terrestres*. São Paulo: Nitro Editorial.
- Tavares, V.C., R. Gregorin and A.L. Peracchi. 2008. A diversidade de morcegos no Brasil: lista atualizada com comentários sobre distribuição e taxonomia; p. 25-60. In S.M. Pacheco, R.V. Marques and C.E.L. Esbérard (org.). *Morcegos do Brasil: biologia, sistemática, ecologia e conservação*. Porto Alegre: Armazém Digital. 575 p.
- Tavares, V.C., L.M.S. Aguiar, F.A. Perini, F.C. Falcão and R. Gregorin. 2010. Bats of the state of Minas Gerais, southeastern Brazil. *Chiroptera Neotropical* 16(1): 675-705.
- Traiano, E. 1985. Ecologia de populações de morcegos cavernícolas em uma região cárscitca do sudeste do Brasil. *Revista Brasileira de Zoologia* 2(5): 255-320.
- Uieda, W., I. Sazima and A. Storti-Filho. 1980. Aspectos da biologia do morcego *Furipterus horrens*. *Revista Brasileira de Biologia* 40(1): 59-66.
- Vieira, C.O. 1942. Ensaio monográfico sobre os quirópteros do Brasil. *Arquivos de Zoologia do Estado de São Paulo* 3(8): 1-471.
- Voss, R.S. and L.H. Emmons. 1996. Mammalian diversity in neotropical lowland rainforests: a preliminary assessment. *Bulletin of the American Museum of Natural History* 230: 1-115.
- Wilson, D.E. 1973. Bat faunas: a trophic comparison. *Systematic Zoology* 22(1): 14-29.

RECEIVED: April 2012

ACCEPTED: October 2012

PUBLISHED ONLINE: December 2012

EDITORIAL RESPONSIBILITY: Valeria da Cunha Tavares